

INDEX.

	Page		Page
<i>Acantharachna mirabilis</i> , <i>Smith</i> , <i>n. sp.</i> (Pl. XVIII.)	335	<i>Ailia affinis</i> , <i>Günther</i> (ftnote)	350
<i>Acanthocystis spinifera</i> , <i>Greeff</i>	294	— <i>coila</i> , <i>Ham. Buch.</i>	350
— <i>turfacea</i>	293	<i>Ailiichthys punctata</i> , <i>Day</i>	350
<i>Acanthonotus Hardwickii</i> , <i>Gray</i> (ftnote)	350	<i>Air-bladder</i> , homologue of	199
<i>Acanthopterygii</i> of India, distrib. of	138, 139	—, opinions respecting	209
<i>Accipiter cirrocephalus</i>	488	<i>Akysis Kurzii</i> , <i>Day</i> (ftnote)	350
<i>Accipitres</i> , geographical distrib. of, 1; of New Guinea	308, 488	<i>Alcyoncellum speciosum</i>	506
<i>Aceros subruficollis</i> , <i>Blyth</i>	156	<i>Alcyone Lessoni</i> , 311, 492; <i>pulchra</i> , 492; <i>pusilla</i>	311, 492
<i>Acilius</i> , 163; <i>fasciatus</i> , breathing exper. on, 180; <i>sulcatus</i>	179	Allman, G. J., recent researches among the more simple sarcode organisms, Anniversary Address for 1876, 261; for 1877	385
<i>Actæa granulata</i>	184	<i>Allorhina anomala</i> , <i>Bates</i>	130
<i>Actæomorpha</i> , <i>Miers</i> , <i>n. gen.</i> , 184; <i>erosa</i> , <i>Miers</i> , <i>n. sp.</i> (Pl. XIV.)	183	<i>Alveolites Battersbyi</i> , <i>Edw. & H.</i> , 360; <i>clavata</i> , 355; <i>compressa</i> , <i>Edw. & H.</i> , 359; <i>cryptodens</i> , <i>Billings</i> , 360; <i>denticulata</i> , <i>Edw. & H.</i> , 360; <i>depressa</i> , <i>Flem.</i> , 359; <i>escharoides</i> , <i>Lam.</i> , 354; <i>Fischeri</i> , <i>Billings</i> , 362; <i>Fougatii</i> , 358; <i>frondosa</i> , <i>Nich.</i> , 362; <i>Goldfussi</i> , <i>Billings</i> , 368; <i>Grayi</i> , <i>Edw. & H.</i> , 359; <i>incrustans</i> , 354; <i>infundibuliformis</i> , 355; <i>Labechii</i> , <i>Edw. & H.</i> , 359; <i>madreporacea</i> , 354; <i>milleporacea</i> , 355; Notes on the genus, 353; <i>repens</i> , <i>Fougat</i> , 360; <i>Rœmeri</i> , <i>Bill.</i> , 369; <i>septosa</i> , <i>Flem.</i> , 357; <i>spongites</i> , <i>Steininger</i> , 355; <i>suborbicularis</i> , <i>Lam.</i> , 354; <i>tubiporacea</i>	355
<i>Actinolophus pedunculatus</i>	303	<i>Ambassia alta</i> , <i>Cuv. & Val.</i> (ftnote), 142; <i>baculis</i> , <i>Ham. Buch.</i> , 143; <i>Barlovi</i> , <i>Sykes</i> (ftnote), 142; <i>Commerstonii</i> , <i>Cuv. & Val.</i> , 143; <i>Dayi</i> , <i>Bleeker</i> , 143; <i>denticulata</i> , <i>Klunz.</i> (ftnote), 143; <i>Dussumieri</i> , <i>Cuv. & Val.</i> (ftnote), 143; <i>gymnocephalus</i> , <i>Lacép.</i> , 143; <i>indica</i> , <i>M. Clell.</i> (ftnote), 142; <i>interrupta</i> , <i>Bleeker</i> , 143; <i>malebaricus</i> , <i>Jerdon</i> (ftnote), 143; <i>nalua</i> ; <i>H. B.</i> , 143; <i>nana</i> , <i>H. B.</i> ,	
<i>Actinometra</i> , 440; and <i>Antedon</i> , the differences, 442; <i>Bennetti</i> , <i>Müll.</i> , 450; <i>cirri</i> , their number, 451; condition of rosette &c., 451; definition of, by Müller, 440; <i>fimbriata</i> , 444; likeness to <i>Solanocrinus</i> , 454; <i>multiradiata</i> , 443; on the genus, 440; <i>polymorpha</i> , morphology of, 440, 444; <i>robusta</i> , 451; <i>solaris</i> , 443; variation in arms &c. of	450		
<i>Actinophrys oculata</i> , 284; <i>sol</i> , 284; <i>viridis</i> , <i>Ehrenb.</i> (ftnote)	292		
<i>Actinosphærium Eichhornii</i>	285, 287		
<i>Actitis hypoleucos</i> (<i>L.</i>)	320, 504		
<i>Æga hirta</i> , 511; <i>macronema</i> , <i>Bleeker</i> , 512; <i>multidigita</i> , <i>Dana</i> , 511; <i>spongiophila</i> , <i>Semper</i> (Pl. XXIV.)	507		
<i>Ægotheles Bennetti</i>	314		
— <i>leucogaster</i>	315		
<i>Ælurædus buccoides</i>	495		
— <i>Stonii</i> , <i>Sharpe</i> , <i>n. sp.</i>	495		
<i>Agabus bipustulatus</i> , exper. on	177		
<i>Ageneiosus childreni</i> , <i>Sykes</i> (ftnote)	349		
<i>Aglyptosternon</i> , <i>Blkr.</i> (ftnote)	343		

	Page		Page
142; oblonga, <i>C. & V.</i> (fnote), 142; ranga, <i>H. B.</i> , 142; thermalis, <i>C. & V.</i> (fnote), 144; Thomassi, <i>Day</i> , 143; urotænia, <i>Bleeker</i> , 143; Vachelli, <i>Peters</i> (fnote)	143	communication, power of, 219; contrast in behaviour of, 238; dependence on slaves, 245; different kinds, 238; different nests, 236; division of labour, 246; experiments testing intelligence, 218; experiments testing the senses, 239; experiments under chloroform and intoxicated, tabular view of, 229; habits of, remarks on, 217; hearing of, 244; individual influence and character, 221; intelligence and affection, 221; intoxicated, 225, 229; parasites of, and on, 257; parthenogenesis in, 257; recollection of friends, 230; routes followed in exp., 240, 245; sight of, 239; slaves of <i>Formica fusca</i> , 248; slaves of <i>Polyergus</i> , 252; suspected cannibalism of, 239; tables of experiments on, 229; tables illustrating experiments on division of labour, 248; want of ingenuity in crossing chasms	217
<i>Amblyceps cæcutiens</i> , <i>Blyth</i> (fn.), 350; mangois, <i>Ham. Buch.</i> , 350; tenuispinus, <i>Blyth</i> (fnote)	350	<i>Aphrodita</i> , sp.	507
<i>Amblyodon</i> (<i>Raf.</i>), (fnote), 140; nicaraguæ, <i>Westw.</i> , = ♂ <i>Phileurus nasicornis</i> , <i>Burm.</i>	130	<i>Apiocrinidæ</i> , new living crinoids belonging to	47
<i>Amblypneustes</i> runs far south, &c.	79	<i>Appendices inferiores</i> in <i>Trichoptera</i> =harpagones	195
<i>Amesia juvenis</i> , 197; pexifascia, <i>Butl.</i> , <i>n. sp.</i> , 115, 197.		<i>Appias plana</i>	197
<i>Amœba actinophora</i> , 438; bilimbosa, 438; polypodia, <i>Max Schultze</i> , 276; princeps, 269; terricola, <i>Greeff</i> , 274 & (nucleus of) 275.		<i>Arcania granulosa</i> , <i>Miers</i> , <i>n. sp.</i>	107
Amœboid changes in plant- and animal-cells	270	<i>Arcella arenaria</i> , 279; hyalina, 399; hyalina, <i>Ehrnb.</i> , = <i>Lecythium</i> , <i>Hert. & Less.</i> , 399; vibrio-like bodies of, 392; vulgaris	279, 391
Amphibious and migratory fishes of Asia	198	rchæopteryx, remark on metacarpus of	101
Amphipnous cuchia, branchial sacs of, 205; dug from clay	214	<i>Archaster andromeda</i> , 71; excavatus, <i>Wy. Thomson</i> , <i>n. sp.</i>	71
Amphistomatous <i>Thalamophora</i>	391	<i>Arcyria flava</i>	414
<i>Amphitrema</i>	410	<i>Ardea cinerea</i> , skull of, 90; flavicollis, <i>Lath.</i> , 504; sacra, <i>Gm.</i>	504
<i>Amphizonella flava</i> = <i>Pseudochlamys patella</i> , 394, 397; vestita, <i>Archer</i> , 277; violacea, <i>Greeff</i> , 276.		<i>Arius pumilus</i> , <i>Val.</i> (fnote), 346; ritoides, <i>Cuv. & Val.</i> (fnote), 346; thalassinus, 338.	
<i>Anabas</i> , <i>Cuv.</i> , 141; kept in pots without water, 201; oligolepis, <i>Günther</i> (fnote), 152; scandens, <i>Dal.</i> , 152; spinosus, <i>Gray & Hardw.</i> (fnote), 152; trifoliatus, <i>Kaup</i> (fnote)	152	<i>Arses Kaupi</i> , 497; telescophtalmus (<i>Garn.</i>), 316, 497.	
<i>Anaitis rosea</i> , <i>McIntosh</i> , <i>n. sp.</i> , description of	214	<i>Arsipoda</i> , <i>Ericks.</i>	477
<i>Anas superciliosa</i> , <i>Gm.</i>	505	— consuta, <i>Germ.</i> , 477; picipes	477
Anderson, Dr. John, on the habits of Hornbills	156	<i>Artamus leucopygialis</i> , <i>Gould</i>	501
<i>Anguilla vulgaris</i> , mode of respiration of	202	<i>Ascaris delphini</i> , 42; maculosa, 45; simplex, 43.	
<i>Anseres</i> of New Guinea	321, 505	Asia, amphibious and migratory fishes of	198
Ant-eater, ear-bones of	188	<i>Aspidobagrus</i> , <i>Bleeker</i> (fnote)	341
<i>Antedon bathyrcinus</i> , 51; celtica, 51; Dubenii, 445, 446 & (calyx of, compared) 456; Eschrichtii, 454; oral pinnules of, irritable, 446; rosacea, 452; rosaceus	446	<i>Asteracanthion</i> , development of	56
<i>Anthias testudineus</i> , <i>Bloch</i> (fnote)	152	<i>Asteridea</i>	71
<i>Anthozoa tabulata</i>	361		
<i>Antichira gratiosa</i> , <i>Sharp</i> , <i>n. sp.</i> , 133; prasina, <i>Burm.</i> , 134; psittacina, <i>Burm.</i>	134		
Ants, 217; chloroformed, 224; com-			

- | | Page | | Page |
|---|--------|---|--------|
| Astictopterus gemmifer, 197; stellerifer, 197. | | Beetles, Australian Phytophagous, 458; description of new species from Central America, 129. | |
| Astrodisculus | 295 | Berardius, ear-bones of | 187 |
| Astur albigularis, 458; approximans, 489; cruentus, 488; Meyerianus, <i>Sharpe, n.sp.</i> (Pl. XXII.), 458; torquatus, 489; Wallacei, 489 | | Bilharzia | 44, 45 |
| Athyma clerica, 196; nivifera | 196 | Birnara nubila | 197 |
| Atta barbara (Pl. XVII.), worker, major and minor | 256 | Boake, Mr., on Ceylon air-breathing fish | 201 |
| Aurelia aurita (Pl. XV.), abnormal forms of, 191; abortion of parts of, 192. | | Bogoda, <i>Bleeker</i> (footnote) | 139 |
| Australian Phytophagous beetles | 458 | Bola, <i>H. B.</i> (footnote) | 140 |
| Bacterium termo | 269 | — chaptis, <i>H. B.</i> (footnote) | 145 |
| Badis, 139, 144; Buchananii, <i>Bleek.</i> , 144; Dario, <i>Ham. Buch.</i> , 144. | | — coitor, <i>H. B.</i> (footnote) | 145 |
| Bagarius Buchananii, <i>Bleeker</i> , 351; Yarrollii, <i>Sykes</i> , 351. | | Bougainvillia Allmanii, <i>Romanes, n. sp.</i> (fnote), 190; fruticosa, name changed to <i>B. Allmanii</i> (footnote) | 190 |
| Bagrus, <i>Cuv. & Val.</i> (fnote) | 341 | Bourguetticrinus, of Chalk and Tertiary formations allied to living Rhizoecrinus | 48 |
| — abbreviatus, <i>C. & V.</i> (fnote) | 344 | Bower Birds of New Guinea | 82 |
| — affinis, <i>Jerdon</i> (fnote) | 345 | Branchiosteus, <i>Gill</i> (footnote) | 342 |
| — agriculus, <i>Jerdon</i> (fnote) | 345 | Brissopsis lyrifera | 67 |
| — albilabrus, <i>C. & V.</i> (fnote) | 344 | British Polyzoa | 479 |
| — aorellus, <i>Blyth</i> (fnote) | 344 | Bucerotidæ of New Guinea | 314 |
| — aorides, <i>Jerdon</i> (fnote) | 344 | Bucharis Chapuisii, 462; granulosus, 462; martius, 463. | |
| — aorinus, <i>Val.</i> (fnote) | 344 | Bucorvus abyssinicus, peculiarities of lachrymal bone of | 91 |
| — Birmanus, <i>C. & V.</i> (fn.) | 344 | Butler, A. G., on new species of Euptychia, with tabular view of those hitherto recorded, 116; on the butterflies of Malacca, 196; on two new Lepidopterous insects from Malacca, 115. | |
| — exodon, <i>Bleeker</i> (fnote) | 345 | Butorides javanica, <i>Horsf.</i> | 320 |
| — fuscus, <i>C. & V.</i> (fnote) | 344 | Butterflies of Malacca | 196 |
| — gulioides, <i>Bleeker</i> (fnote) | 344 | —, geographical distribution of | 197 |
| — keletius, <i>Bleeker</i> (fnote) | 345 | Cacomantis assimilis, <i>Gray</i> | 492 |
| — Lamarrii, <i>C. & V.</i> (fnote) | 344 | Callichrous affinis, <i>Swains.</i> (fnote) | 348 |
| — melas, <i>Bleeker</i> (fnote) | 344 | 348; bimaculatus, <i>Bloch</i> , 348; ceylonensis, <i>Günth.</i> (fnote), 348; Egertonii, <i>Day</i> (fnote), 348; gangeticus, <i>Peters</i> , 348; immaculatus, <i>Swains.</i> (fnote), 348; macrophthalmus, <i>Blyth</i> , 348; macrostomus, <i>Swains.</i> (fnote), 348; malabaricus, <i>C. & V.</i> , 348; nebulosus, <i>Swains.</i> (fnote), 348; nigrescens, <i>Day</i> (fnote), 348; notatus, <i>Day</i> (fnote), 348; pabda, <i>Ham. Buch.</i> , 348; pabo, <i>H. B.</i> , 348; sindensis, <i>Day</i> , 348; vittatus, <i>Swains.</i> (fnote), 348. | |
| — rhodopterygius, <i>Blk.</i> (fnote) | 344 | Callidula abisara | 197 |
| — Schlegelii, <i>Blk.</i> (fnote.) | 344 | Calliplœa leucogonis | 196 |
| — tengora, var., <i>Blyth</i> (fnote) | 345 | Callomystax, <i>Günth.</i> (fnote) | 343 |
| — trachacanthus, <i>C. & V.</i> (fn.) | 345 | Calomela, <i>Hope</i> | 475 |
| Bairdiella, <i>Gill</i> (fnote) | 140 | Calornis cantoroides | 501 |
| Balæna, ear-bones of | 187 | | |
| Balæniceps rex, orbital circle of | 91 | | |
| Balænoptera, malleus and incus of | 187 | | |
| Baly, J. S., on species of Australian Phytophagous beetles | 458 | | |
| Barbel, exper. on its respiration | 203 | | |
| Barbus | 202 | | |
| Batassio, <i>Blyth</i> (fnote), 343; affinis, <i>Blyth</i> (fnote), 344. | | | |
| Bathybius, remarks as to its nature | 433 | | |
| Bathycrinus Aldrichianus, <i>Wy. Thomson, n. sp.</i> , 47, 49, 50, 54; gracilis, <i>Wy. Thom., n. sp.</i> | 47, 54 | | |
| Baza Reinwardti, <i>Müll. & Schl.</i> , 309, 490; suberistata, 490. | | | |
| Bedula, <i>Gray</i> (fnote) | 139 | | |
| — Hamiltonii, <i>Gray & Hardw.</i> (fnote), 144. | | | |

- | | Page | | Page |
|--|----------|---|----------|
| Calornis mysolensis | 501 | Chanos salmoneus, introd. into tank | 211 |
| — viridescens | 318, 501 | Charadrius fulvus, <i>Gm.</i> | 504 |
| Calycine monad, development of . | 423 | Cheilodipterus culius, <i>H. B.</i> (ftnote) | 148 |
| Campula oblonga, <i>Cobbold</i> , = | | Cheilotrema, <i>v. Tsch.</i> (ftnote) . . | 140 |
| Distoma campula | 40 | Chibia assimilis | 500 |
| Canda reptans (Pl. XXIII.) | 486 | — carbonaria (<i>Müll.</i>) | 499 |
| Caprimulgus macrurus | 314, 493 | Chimarichthys, <i>Sauv.</i> (ftnote) . . | 344 |
| Carnivorous Water-beetles, respira- | | Chirodota, larva of, taken by 'Chal- | |
| tory action of | 161 | lenger' expedition | 79 |
| Carpenter, P. Herbert, on Actino- | | Chiroptera, ossicula auditus of . . | 186 |
| metra, <i>Müll.</i> , with account of a | | Chlamidomyxis labyrinthuloides, | |
| new species from the Philippines | 440 | <i>Archer</i> | 281, 282 |
| Carpophaga muelleri, 319, 502; | | Chlamydodera cerviniventris, <i>Gould</i> | |
| pinon, 319, 502; puella, 503; | | 82, 495 | |
| zoæ, 502. | | Chlamydophrys stercorea, <i>Cienk.</i> , | |
| Carpus buried in mud (ftnote) . . | 215 | 399; probably = <i>Diffugia enche-</i> | |
| Carystea, <i>Baly</i> | 475 | lyls, <i>Schneid.</i> , 400. | |
| Castalius ethion | 196 | Chaetetes depressa, <i>Flem.</i> , 367; hy- | |
| Catallacta | 438 | perboreus, <i>Nich. & Eth.</i> , 367; ra- | |
| Catargyra elegans | 197 | dians, <i>Fischer</i> , 365; septosus, | |
| Catharistes atratus, geog. distrib. of, | | <i>Flem.</i> , 367. | |
| 19; remark on, 23; map show- | | Chaetodon kakaitzel, <i>Lacép.</i> (ftnote) | 1 & 3 |
| ing range of, Pl. III. | | Chromides | 141, 153 |
| — aura, remark on | 23 | Chrysochloris, ossicula auditus of . | 186 |
| Cathartes papa, geog. distrib. of, 18; | | Chrysomelidæ | 474 |
| map showing range of, Pl. II. | | Cidaris abyssicola, 64; affinis (<i>Sto-</i> | |
| Catopra, <i>Bleeker</i> (ftnote) | 139 | kesii), 64; hystrix, 64; nutrix, | |
| — siamensis, <i>Günth.</i> (ftnote) . . . | 144 | <i>Wy. Thoms.</i> , <i>sp. n.</i> , 62, 63; pa- | |
| — tetracanthus, <i>Günth.</i> (ftnote) . . | 144 | pillata, 62, 64; papillata, var. | |
| Catylopus, <i>Guich.</i> (footnote) . . . | 140 | <i>Stokesii</i> , 63. | |
| Centropus bernsteinii, 81; mene- | | Ciliophrys infusionum | 289 |
| beki, 81; 311, 491; spilopterus, | | Cinnyris frenatus, <i>S. Müll.</i> | 316, 496 |
| 81, 310, 491; violaceus, 81. | | Cirolana multidigitata (Pl. XXIV.) | 511 |
| Ceratodus, air-bladder converted | | Cirrochroa rotundata | 196 |
| into a lung in | 198 | Cisticola ruficeps | 315, 494 |
| Ceryle maxima, skull of | 90 | Cladodaetyla crocea, <i>Less.</i> | 57, 58 |
| Cetacea, ossicula auditus of | 187 | Clarias assamensis, <i>Day</i> , 349; bra- | |
| Cethosia methypsea | 196 | chrysoma, <i>Günth.</i> (ftnote), 349; | |
| Ceylon fishes, note on | 139 | Dussumieri, <i>Cuv. & Val.</i> , 349; | |
| Ceyx solitaria, <i>Temm.</i> | 492 | dendritic branchial cavity of, 205; | |
| Chaca Buchanani, <i>Günth.</i> (ftnote) | 349 | magur, <i>Ham. Buch.</i> 349; marpus, | |
| — lophioides, <i>Cuv. & Val.</i> | 349 | <i>Cuv. & Val.</i> (ftnote), 349; me- | |
| Chalarothoraca | 298 | lanosoma, <i>Bleek.</i> (ftnote), 349; | |
| Chalcophaps chrysochlora (<i>Wagl.</i>), | | punctatus, <i>Cuv. & Val.</i> (ftnote), | |
| 319, 502; stephani, <i>P. & Jacq.</i> | 502 | 349; <i>Teysmanni</i> , <i>Bleek.</i> , 349. | |
| Chalcopsittacus chloropterus, | | Clarke, R., and Mivart, St. G., on | |
| <i>Salvad.</i> | 491 | sacral plexus and vertebræ of | |
| 'Challenger' Expedition, Sir C. | | Lizards | 370 |
| Wy. Thomson on Crinoids ob- | | Clathrulina elegans | 299 |
| tained in the | 47 | Clupisoma argentata, <i>Swains.</i> (foot- | |
| Channa, <i>Gronov.</i> | 141 | note) | 342, 347 |
| — indica, <i>Gronov.</i> (ftnote) | 152 | Cobalus ciliatus | 197 |
| — orientalis, <i>Bl. Schn.</i> | 152 | Cobbold, Dr. T., Trematode para- | |
| Chamaëlo vulgaris, sacral plexus of | 371 | sites from Dolphins of Ganges . . | 35 |
| Chanda, <i>Ham. Buch.</i> (ftnote), 139; | | Cobitis pacifica, <i>Forst.</i> (ftnote) . . | 148 |
| bagoda, <i>H. B.</i> (footnote), 142; | | Coccospheres, nature of | 484 |
| lala, <i>H. B.</i> (ftnote), 142; phula, | | Cochliopodium pellucidum | 277 |
| <i>H. B.</i> (ftnote), 142; roconius, | | Cœnites orientalis, <i>Eichw.</i> | 361 |
| <i>M' Clell.</i> (ftnote), 142. | | Cojus cobojus, <i>Ham. Buch.</i> (ftnote) | 152 |

- | | Page | | Page |
|---|----------|---|------|
| Colisa, <i>Cuv. & Val.</i> (fnote) . . . | 141 | Ctenops, <i>M' Clell.</i> (fnote) . . . | 141 |
| — ponticeriana, <i>C. & V.</i> (fnote) . | 153 | Cuchia Eel, branchial arches of . . | 205 |
| — unicolor, <i>C. & V.</i> (fnote) . . . | 153 | Cyathocerinus | 53 |
| Collett, Robert, on <i>Myodes lemmus</i> | | Cybister <i>Rosseli</i> | 163 |
| in Norway | 327 | Cyclocephala <i>conspicua</i> , <i>Sharp</i> , | |
| Collocalia terræ-reginæ, <i>Ramsay</i> . | 494 | <i>n. sp.</i> | 135 |
| Collosphæra, Cienkowski on . . . | 426 | — <i>proba</i> , <i>Sharp, n. sp.</i> | 135 |
| Colluricincla brunnea, <i>Gould</i> . . . | 499 | Cyclonoda, <i>Baly, n. g.</i> , 474; <i>pilula</i> , | |
| — pallidirostris, <i>Sharpe</i> | 499 | <i>Clark</i> , 474; <i>subpunctata</i> , <i>Clark</i> . | 474 |
| Colpodella, reproduction of . . . | 423 | Cyclopsitta <i>suavissima</i> , <i>Sclater</i> . | 491 |
| Columbæ of New Guinea | 318, 502 | Cyclopsittacus <i>cervicalis</i> | 310 |
| Colymbetes exoletus, exper. on . . | 175 | Cyclothurus, malleus of | 188 |
| Comaster | 455 | Cynorhamphus, neck-vertebræ of . | 94 |
| Comatula (Alecto) and Actinometra, | | — <i>suævicus</i> , palatal bones of . . . | 92 |
| distinctions of, 442, 451; Müller's | | Cynthia erotella | 196 |
| classification of, artificial, 441; | | Cyphoderia <i>margaritacea</i> , <i>Schumb.</i> | 406 |
| multiradiata, 441; polymorpha, | | — <i>truncata</i> , <i>Schulze</i> | 407 |
| 441; rosea, 442; trichoptera, | | Cypselus <i>pacificus</i> (<i>Lath.</i>) | 494 |
| 441; what comes under, 441. | | Cystophrys <i>Hæckeliana</i> , <i>Archer</i> . | 402 |
| Condylura, malleus and incus of . . | 186 | — <i>oculea</i> , nature of | 409 |
| Coracinus, <i>Pall.</i> (fnote) | 140 | Cytode | 262 |
| Corals, Palæozoic | 353 | Dacelo <i>gaudichaudi</i> , 313, 493; in- | |
| Cornuspira, Max <i>Schultze</i> on . . . | 262 | <i>intermedius</i> , <i>Salvad.</i> , 312, 493. | |
| Corone orru, <i>Müll.</i> | 318 | Dactylosphærium <i>vitreum</i> | 266 |
| Corvina, <i>Cuv.</i> (fnote), 140; cele- | | Dasybus, <i>ossicula auditus</i> of . . . | 188 |
| bica, <i>Bleek.</i> (fnote), 145; dor- | | Day, Francis, geographical distribu- | |
| salis, <i>Peters</i> (fnote), 145; <i>Kuhlii</i> , | | of Indian Freshwater Fishes. | |
| <i>Cuv. & Val.</i> (fnote), 145; <i>lobata</i> , | | — Part I. The Acanthopterygii, | |
| <i>Cuv. & Val.</i> (fnote), 145; <i>Neillii</i> , | | Spiny-rayed Teleostean Fishes . | 138 |
| <i>Day</i> (fnote), 145; <i>platycephala</i> , | | — Part II. The Siluridæ | 338 |
| <i>Cuv. & Val.</i> (fnote), 145; <i>sampitensis</i> , | | — on amphibious and migratory | |
| <i>Bleek.</i> (fnote), 145; <i>soldado</i> , | | Fishes of Asia | 198 |
| <i>Cant.</i> (fnote), 145; <i>Wolfii</i> , | | — on some Irish Gasterostei . . | 110 |
| <i>Bleeker</i> (fnote), 145. | | Defrancia <i>lucernaria</i> | 485 |
| Corvus orru, <i>Müll.</i> | 501 | Delias <i>metarex</i> | 197 |
| Corymbipora <i>fungiformis</i> , <i>Smitt</i> . | 485 | Delphinus, ear-bones of | 187 |
| Cossyphus, <i>M' Clell.</i> (fnote) . . . | 343 | Dendrocygna <i>guttata</i> , 505; <i>vagans</i> , | 505 |
| Cotinis <i>adpersa</i> , <i>Sharp, n. sp.</i> . . . | 136 | Description of new species of Beetles | |
| — <i>gracilis</i> , <i>Sharp, n. sp.</i> | 157 | from Central America | 129 |
| Cracticus <i>cassicus</i> (<i>Bodd.</i>), 82, 317, | | Desmothoraca | 298 |
| 499; <i>mentalis</i> , <i>Salvad. & D'Al-</i> | | Diadema <i>incommoda</i> | 196 |
| <i>bert.</i> , 317, 499; <i>quoyi</i> , <i>Less.</i> , 499. | | Dialithus <i>magnificus</i> | 130 |
| Craniorrhinus <i>cassidix</i> , 156; <i>wal-</i> | | Diaphoropodon <i>mobile</i> , <i>Archer</i> . | 408 |
| <i>deni</i> , <i>Sharpe, n. sp.</i> | 156 | Dicaëum <i>rubrocoronatum</i> , <i>Sharpe</i> . | 496 |
| Crematogaster <i>scutellaris</i> | 223 | Dichocerinus | 51 |
| Crinoids, notice of new living, 47; | | Didosaurus, <i>Günth., n. gen.</i> , 322; | |
| subepithelial band, nature of . . | 447 | <i>mauritanus</i> , <i>Günth., n. sp.</i> , (bones | |
| Crotch, W. Duppa, on the migra- | | of) 322 & (affinities of) 324. | |
| tion and habits of Norwegian | | Didymium <i>serpula</i> | 414 |
| Lemming, 27; additional note | | Diffugia <i>acropodia</i> , <i>Hertw. & Less.</i> , | |
| on, 83; further remarks on, 157. | | 396; <i>enclélus</i> , 398; <i>symmetrica</i> , | |
| Crustacea, new genus and species | | <i>Wallich</i> , 395. | |
| of, 183; <i>oxystomatous</i> , E. J. | | Dimorphodon, 98; <i>vertebræ</i> of, 94. | |
| Miers on, 107; species of, living | | Diphyllbothrum <i>stemmaccephalum</i> | |
| within Venus's Flower-basket . . | 506 | from Porpoise | 42 |
| Cryptocephalidæ | 459 | Diplolepis, <i>Steind.</i> (fnote) | 140 |
| Cryptocnemius <i>holdsworthi</i> , <i>Miers</i> , | | Diplophrys <i>Archeri</i> | 408 |
| <i>sp. nov.</i> | 107 | Discoliths, nature of | 433 |
| Cryptopterus <i>latovittatus</i> , <i>Pl.</i> (fn.) | 348 | | |

- | | Page | | Page |
|--|------|---|------|
| <i>Discopora hispida</i> , var. <i>meandrina</i> , 483; <i>meandrina</i> , <i>Peach</i> , <i>n. sp.</i> Pl. XXIII., 482. | | 148; <i>macrolepidota</i> , <i>Bloch</i> , 148; <i>margaritacea</i> , <i>Cuv. & Val.</i> (ftnote), 148; <i>mauritanicus</i> , <i>Benn.</i> (ftnote), 148; <i>muralis</i> , <i>Cuv. & Val.</i> , 148; <i>nigra</i> , <i>Quoy & Gaim.</i> (ftnote), 148; <i>ophiocephalus</i> , <i>Cuv. & Val.</i> , 148; <i>porocephalus</i> , <i>Cuv. & Val.</i> , 148; <i>prismatica</i> , <i>Bleeker</i> (footnote), 149; <i>scintillans</i> , <i>Blyth</i> , 148; <i>sexguttata</i> , <i>Cuv. & Val.</i> , 148; <i>soaresi</i> , <i>Playfair</i> (ftnote), 148; <i>viridis</i> , <i>Bleeker</i> (ftnote), 148. | |
| <i>Distoma Andersoni</i> , <i>Cobbold</i> , <i>n. sp.</i> , 46; <i>campula</i> , <i>Cob.</i> , 40 & (anatomy of) 41; <i>compactum</i> from an <i>Ichneumon</i> , 41; <i>crassum</i> , uterine folds of ♀, 40; <i>lancea</i> , <i>Diesing</i> , 36 & (anatomy of) 37. | | <i>Embia Batesii</i> , <i>M' Lach.</i> , <i>sp. n.</i> , 380; <i>Klugi</i> , <i>Ramb.</i> , 380; <i>Latreillii</i> , 379; <i>mauritanica</i> , 375; <i>nigra</i> , <i>Hagen</i> , 383; <i>persica</i> , <i>M' Lach.</i> , <i>sp. n.</i> , 382; <i>Salvini</i> , <i>M' Lach.</i> , <i>sp. n.</i> , 380; <i>Savignyii</i> , 376; <i>Solieri</i> , <i>Ramb.</i> , 376. | |
| <i>Ditropidus costipennis</i> , <i>Baly</i> , <i>n. sp.</i> , 465; <i>facialis</i> , <i>Baly</i> , <i>n. sp.</i> , 466; <i>Jansoni</i> , <i>Baly</i> , <i>n. sp.</i> , 467; <i>lætus</i> , <i>Baly</i> , <i>n. sp.</i> , 464; <i>ornatus</i> , <i>Baly</i> , <i>n. sp.</i> , 467; <i>phalacroides</i> , <i>Baly</i> , <i>n. sp.</i> , 464; <i>pulchellus</i> , <i>Baly</i> , <i>n. sp.</i> , 468; <i>serenus</i> , <i>Baly</i> , <i>n. sp.</i> , 468. | | <i>Embidiæ</i> , distribution of family, 379; habits of the family, 375; historical remarks on, 374; number of species, 379; on the nymph-stage of, 373; systematic position, structure, &c., 377. | |
| Dolphins, ear-bones of 187 | | Encysting process, <i>Hertwig & Lesser</i> on 265 | |
| <i>Domicella scintillata</i> , <i>Temm.</i> 80 | | <i>Entosolenia globosa</i> , nucleus in 411 | |
| <i>Domopora</i> (<i>Defrancia</i>) <i>stellata</i> , <i>Goldf.</i> , 484; <i>truncata</i> , <i>Jameson</i> , 483. | | <i>Epinephile hyperantus</i> , <i>L.</i> 195 | |
| <i>Donacola castaneothorax</i> , 502; <i>nigriceps</i> , <i>Ramsay</i> , 501. | | <i>Erethistes conta</i> , <i>Ham. Buch.</i> , 346; <i>elongata</i> , <i>Day</i> , 346; <i>hara</i> , <i>Ham. Buch.</i> , 346; <i>Jerdonii</i> , <i>Day</i> , 346; <i>pusillus</i> , <i>Müll. & Trosch.</i> (ftnote) 346. | |
| <i>Doran</i> , <i>Alban H. G.</i> , morphology of mammalian ossicula auditus . 185 | | <i>Eschara cervicornis</i> (ftnote), 482; <i>rosacea</i> , 481; <i>Skenci</i> , var. <i>tridentata</i> , <i>Busk</i> (Pl. XXIII.), 480; <i>stellata</i> , <i>Peach</i> , <i>n. sp.</i> (Pl. XXIII.), 481. | |
| <i>Dysectodes</i> 263 | | <i>Etheridge</i> , <i>Robert</i> , and <i>Nicholson</i> , <i>H. Alleyne</i> , on the genus <i>Alveolites</i> 353 | |
| <i>Dytiscidæ</i> , on the respiratory action of 161 | | <i>Etroplus</i> , <i>Cuv. & Val.</i> , 141; <i>canarensis</i> , <i>Day</i> , 153; <i>coruchi</i> , <i>Cuv. & Val.</i> (ftnote), 153; <i>maculatus</i> , <i>Bloch</i> , 153; <i>meleagris</i> , <i>Cuv. & Val.</i> (ftnote), 153; <i>suratensis</i> , <i>Bloch</i> , 153. | |
| <i>Dytiscus marginatus</i> , 161; breathing-orifices of, 162, of ♂, 181, of ♀, 182. | | <i>Eudynamis cyanocephala</i> 310, 491 | |
| Ear-bones, small, of mammals, abstr. of paper on 185 | | <i>Euglypha alveolata</i> , 398; <i>ampulacea</i> , <i>Dujard.</i> , 404; <i>pleurostoma</i> , <i>Carter</i> , 398. | |
| <i>Echidna</i> , ankylosis of malleus and incus of 189 | | <i>Euglyptosternum lineatum</i> , <i>Day</i> . 352 | |
| <i>Echinaster</i> , development of 56 | | <i>Eulabes dumonti</i> (<i>Less.</i>) . 82, 318, 501 | |
| <i>Echinoderms</i> , peculiarities in propagation of, 55; southern, rear young in a pouch 56 | | <i>Eunolpidæ</i> 469 | |
| <i>Echinoidea</i> , <i>demosticha</i> , 62; <i>petalosticha</i> , 67. | | <i>Euphoria Belti</i> , <i>Sharp</i> , <i>n. sp.</i> . . . 137 | |
| <i>Echinus</i> , runs far south 79 | | <i>Euplectella aspergillum</i> , 506; <i>Crustacea</i> living within, 506; <i>speciosa</i> 506 | |
| <i>Eclectus polychlorus</i> (<i>Scop.</i>) 491 | | <i>Euplœa Pinwillii</i> 196 | |
| <i>Edentata</i> , ear-bones of 188 | | | |
| <i>Edoliosoma melas</i> 317 | | | |
| <i>Eleotris</i> , <i>Gronov.</i> , 141; <i>amboinensis</i> , <i>Bleek.</i> , 149; <i>baccata</i> , <i>Blyth</i> (ftnote), 149; <i>butis</i> , <i>Ham. Buch.</i> , 149; <i>canariensis</i> , <i>Day</i> , 148; <i>cantoris</i> , <i>Günth.</i> (ftnote), 148; <i>caperata</i> , <i>Cantor</i> , 149; <i>cavifrons</i> , <i>Blyth</i> , 148; <i>feliceps</i> , <i>Blyth</i> , 148; <i>fusca</i> , <i>Bl. Schn.</i> , 148; <i>koilomatodon</i> , <i>Bleeker</i> (ftnote), 149; <i>humeralis</i> , <i>Cuv. & Val.</i> (ftnote), 149; <i>incerta</i> , <i>Blyth</i> (ftnote), 148; <i>litoralis</i> , <i>Day</i> , 148; <i>lutca</i> , <i>Day</i> , | | | |

Page

Euptychia abretia, 117; *albofasciata*, 117; *acemnis*, 119; *ægrotata*, 121; *ætherialis*, *Butl., n. sp.* (Pl. XII.), 126; *affinis*, 118; *agatha*, 122; *alcinoë*, 123; *ambigua*, 119; *ambigua group*, 119; *anacleta*, *Butl., n. sp.* (Pl. XII.), 123; *angelica*, 121; *angularis*, 118; *antonoë*, 120; *argante*, 118; *argentella*, 122; *argyrospila*, 118; *armilla*, 117; *arnæa*, 122; *arnæa group*, 122; *ashna*, 117; *atalanta*, 119; *austeria*, 118; *ayaya*, 122; *Batesii*, 121; *benedicta*, *Butl., n. sp.* (Pl. XII.), 124; *binalinea*, 119; *binocula*, 117; *brixiola*, 121; *brixius*, 121; *byses*, 122; *cærulea*, 121; *calixta*, *Butl., n. sp.* (Pl. XII.), 125; *callichloris*, 122; *calpurnia*, 116; *camerta*, 119; *celmis*, 118; *chloris*, 122; *elementia*, *Butl., n. sp.* (Pl. XII.), 128; *clorimena*, 122; *clorimena group*, 122; *cluena*, 121; *cluena group*, 122; *cœlestis*, 121; *cœlica*, 121; *cornelius*, 122; *cosmophila*, 122; *crantor*, 118; *cyanites*, 121; *cyclops*, *Butl., n. sp.* (Pl. XII.), 126; *disaffecta*, 118; *divergens*, 118; *doris*, 121; *doris group*, 121; *doxes*, 122; *doxes group*, 122; *electra*, 118; *enyo*, 117; *ecüs*, 119; *epinephele*, 119; *erichtho*, 121; *erigone*, 118; *eriphule*, 118; *erycina*, 122; *eurytus*, 118; *eusebia*, *Butl., n. sp.* (Pl. XII.), 126; *fabiana*, *Butl., n. sp.* (Pl. XII.), 126; *fallax*, 119; *fetna*, 117; *francisca*, 117; *fulgora*, 121; *fumata*, 120; *furnia*, 117; *galesus*, 119; *gemmula*, 122; *gera*, 121; *gera group*, 121; *gigas*, 120; *glauцина*, 121; *grimon*, 119; *grimon group*, 119; *gulnari*, 120; *harmonia*, 120; *harmonia group*, 120; *helle*, 117; *hermes*, 119; *hermes group*, 119; *herse*, 122; *hewitsonii*, 122; *hiemalis*, 121; *hilara*, 117; *Hübneri*, 119; *hygnia*, *Butl.* (Pl. XII.), *n. sp.*, 127; *insignis*, 121; *insolata*, 117; *iris*, 122; *ithama*, 123; *itonis*, 122; *jesia*, 117; *jovita*, 119; *junia*, 121; *junonia*, 122; *labè*, 117; *languida*, 116; *latia*, 122; *lea*, 121; *lea group*, 121; *lethe*, 118; *libye*, 120; *libye group*, 120; *libyoidea*, 120; *litrata*, 117;

Page

litrata group, 117; *lobelia*, 121; *lupita*, 123; *lydia*, 116; *macrophthalma*, 117; *mæpius*, 122; *mæpius group*, 122; *mainmouné*, 119; *manasses*, 123; *mare*, 121; *marmorata*, 119; *melchiades*, *Butl., n. sp.* (Pl. XII.), 125; *melobosis*, 119; *metagera*, 121; *metaleuca*, 116; *mina*, 120; *modesta*, 119; *mollina*, 117; *mollina group*, 117; *mollis*, 117; *muscosa*, 120; *myncea*, 118; *nana*, *Mæsch., n. sp.*? a var. of *E. hermes*, 128; *neecys*, 120; *neecys group*, 120; *nortia*, 121; *nossis*, 117; *nossis group*, 117; *numeraria*, 123; *numilia*, 123; *obscura*, 120; *ochracea*, 118; *ocirrhoë*, 116; *ocirrhoë group*, 116; *oenus*, 118; *ocypete*, 117; *ocypete group*, 117; *oreba*, 120; *pacarus*, 120; *pacarus group*, 120; *pæon*, 118; *pagyris*, 122; *palladia*, 118; *peculiaris*, 121; *peloria*, 123; *penelope*, 118; *periphass*, 118; *pharella*, 119; *phares*, 119; *phares group*, 119; *philippa*, 121; *phineas*, 120; *phocion*, 122; *phocion group*, 122; *phronius*, 118; *picea*, 122; *peria*, 118; *pilata*, 121; *pimpla*, 119; *poltys*, 119; *polyphemus*, 120; *pompilia*, 123; *pronophila*, 117; *pronophila group*, 117; *pyracmon*, 122; *quadrina*, 122; *quantus*, 120; *quantus group*, 120; *renata*, 118; *renata group*, 118; *rubricata*, 119; *rustica*, 120; *sabina*, 120; *salvini*, 122; *saturnus*, 117; *satyrina*, 120; *Saundersii*, 120; *sericella*, 122; *similis*, 118; *sosybius*, 119; *soter*, *Butl., n. sp.* (Pl. XII.), 124; *species?*, 116; *stelligera*, 120; *straminea*, 118; *sylvina*, 123; *squamistriga*, 120; *telesphora*, *Butl., n. sp.* (Pl. XII.), 127; *terentia*, 117; *terrestris*, 118; *thalesa*, *Mæsch., n. sp.*? an abnormal form of *E. batesii*, 128; *themis*, *Butl.*, 118; *tiessa*, 120; *tolumnia*, 122; *tricolor*, 121; *umbrosa*, 119; *undetermined species of*, 123; *undina*, 118; *undulata*, 119; *urania*, 121; *urbana*, *Butl., n. sp.* (Pl. XII.), 124; *usitata*, 118; *variabilis*, 118; *vesper*, 117; *vesta*, 117; *vestigata*, 118; *west-*

- | | Page | | Page |
|---|---------|--|----------|
| woodii, 117; ypthima, 120; | | Geckoid Lizard, bones of, from | |
| zeba, 120; zeredatha, 119; ziza | 121 | Mascarene Islands | 324 |
| European Rhopalocera, on the male | | Geloptera igneo-nitens, <i>Baly</i> , <i>n. sp.</i> | 472 |
| genital armature of | 195 | — vestita, <i>Baly</i> , <i>n. sp.</i> | 473 |
| Eurystomus crassirostris, <i>Sclater</i> | 493 | Genital armature of European Rho- | |
| Eutropius macrophthalmus, <i>Blyth</i> | | palocera | 195 |
| (ftnote) | 347 | Genyonemus, <i>Gill</i> (ftnote) | 140 |
| Eutropiichthys Burmanicus, 350; | | Geoffroyus aruensis | 309, 491 |
| vacha, <i>Ham. Buch.</i> , 350. | | — pucherani | 80 |
| Exostoma Berdmorei, <i>Blyth</i> , 352; | | Geographical distribution of Acci- | |
| Blythii, <i>Day</i> , 352; labiatum, | | pitres, 1; of Indian freshwater | |
| <i>M'Clell.</i> , 352; Stoliczkæ, <i>Day</i> , 352 | | fishes, Part. I., 138; Part II., | |
| Experiments on air-breathing fishes | 203 | 338; of Vulturidæ, 1. | |
| — on breathing of water- | | Geopelia humeralis (<i>Temm.</i>) | 503 |
| beetles | 162-183 | Georychus probably a Lemming | 34 |
| Extinct Mascarene Lizards | 322 | Gerygone chrysogastra, <i>Gray</i> | 495 |
| | | — cinerascens, <i>Sharpe</i> , <i>n. sp.</i> | 494 |
| Fasciola Jacksoni | 43 | — palpebrosa | 495 |
| Faula centralis, <i>Sharp</i> , <i>n. sp.</i> | 131 | Globiocephalus, ear-bones of | 187 |
| Filifera of Schulze = Reticularia, | | Glyciophila modesta, <i>Gray</i> , 316, | |
| <i>Carpenter</i> | 390 | 496; subfasciata, <i>Ramsay</i> , 496. | |
| — rich in species | 396 | Glyphodes amethysta | 197 |
| Fish, abnormal variations of, 110; | | Glyptosternum botia, <i>Ham. Buch.</i> , | |
| amphibious and migratory of | | 351; cavia, <i>H. B.</i> , 351; con- | |
| Asia, 198; exhumation from | | rostre, <i>Steind.</i> , 351; dekkanense, | |
| earth, 214; falling from clouds, | | <i>Günth.</i> (ftnote), 351; gracile, | |
| 213; poisoned by muddy water, | | <i>Günth.</i> (ftnote), 351; lonah, | |
| 200. | | <i>Sykes</i> , 351; madraspatanum, | |
| Fluke, from biliary ducts of Por- | | <i>Day</i> , 351; modestum, <i>Day</i> | |
| poise, 38; from Giraffe, 43; | | (ftnote), 351; pectinopterum, | |
| larvæ of Elephant, 43. | | <i>M'Clell.</i> 351; Stoliczkæ, <i>Steind.</i> | |
| Foraminifera, nucleus in | 410 | (ftnote), 351; striatum, <i>M'Clell.</i> , | |
| Formica cinerea, 257; fusca (Pl. | | 351; telchitta, <i>H. B.</i> , 351; | |
| XVII.), 223, 230, 237, 246; | | trilineatum, <i>Blyth</i> , 351. | |
| nigra, 218, 221; sanguinea (Pl. | | Glyptothorax, <i>Blyth</i> (ftnote) | 343 |
| XVII.), 256. | | Gobiidæ | 140 |
| Freshwater fishes of India, geog. | | Gobius acutipennis, <i>Cuv. & Val.</i> , | |
| distrib. of | 138 | 147; albo-punctatus, <i>C. & V.</i> , | |
| | | 147; andamanensis, <i>Day</i> (ftnote), | |
| Gagata batasio, <i>Ham. Buch.</i> , 350; | | 146; artedi, 140; biocellatus, | |
| cenia, <i>H. B.</i> , 350; itchkeea, | | <i>C. & V.</i> , 147; Bleekeri, <i>Day</i> , | |
| <i>Sykes</i> , 350; tengana, <i>H. B.</i> , 350. | | 147; breviceps, <i>Blyth</i> (ftnote), | |
| Gagata typus, <i>Bleek.</i> (ftnote) | 350 | 147; brevifilis, <i>C. & V.</i> (ftnote), | |
| Galeopithecus, ossicula of | 186 | 146; brevirostris, <i>Günth.</i> , 146; | |
| Gallurucidæ | 475 | bullee-kokah, <i>Russ.</i> (ftnote), | |
| Gallinula ruficrissa, <i>Gould</i> | 505 | 146; bynoensis, <i>Richardson</i> , 146; | |
| Gangetic dolphins, parasites of | 35 | caninus, <i>Günth.</i> (ftnote), 146; | |
| Gasterosteï, on some Irish | 110 | catebus, <i>C. & V.</i> (ftnote), 146; | |
| Gasterosteidæ, remarks on Dr. Sau- | | celebicus, <i>C. & V.</i> (ftnote), 147; | |
| vage's revision of | 110 | chlorostigma, <i>Bleek.</i> (ftnote), | |
| Gasterosteus aculeatus, <i>Linn.</i> , 113, | | 146; criniger, <i>C. & V.</i> , 146; cris- | |
| variations of, 114; pungitius, | | tatus, <i>Day</i> , 147; cyanomos, <i>Blk.</i> , | |
| <i>Linn.</i> , 112, red coloration of | | 146; elegans, <i>C. & V.</i> , 147; | |
| (ftnote), 111, variations of, 113; | | giuris, <i>Ham. Buch.</i> , 146; giuris | |
| spinachia, 112. | | captured at sea, 211; gobiodon, | |
| Gecko Newtonii, <i>Günth.</i> , <i>n. sp.</i> , | | <i>Day</i> (ftnote), 147; grammepo- | |
| 325; description, bones of, 325; | | mus, <i>Bleek.</i> (ftnote), 147; gri- | |
| verus, 324, 326. | | seus, <i>Day</i> , 146; gutum, <i>H. B.</i> , | |

Page

Page

- 147; interstinctus, *Rich.* (fnote), 147; kokius, *C. & V.* (fnote), 146; koku, *Russ.* (fnote), 146; kora, *C. & V.* (fnote), 146; korah mottah, *Russ.* (fnote), 146; Kreffii, *Steind.* (fnote), 146; kurpah, *Sykes* (fnote), 146; litturatus, *Steind.* (fnote), 147; macrostoma, *Steind.*, 146; macrurus, *Bleek.* (fnote), 147; madraspatensis, *Day*, 147; magniloquus, *Day*, 147; malabaricus, *Day*, 147; Masoni, *Day*, 146; melanocephalus, *Bleek.* (fnote), 147; melanosoma, *Bleek.*, 147; melanosticta, *Day*, 147; Neilli, *Day*, 147; novemradiatus, *Ham. Buch.* (fnote), 148; nuna mottah, *Russ.* (fnote), 146; nunus, *H. B.*, 147; ocellatus, *Day*, 146; ornatus, *Rüpp.*, 147; padangensis, *Bleek.* (fnote), 147; periophthalmoides, *Blk.* (fnote), 147; personatus, *Bleek.*, 147; planiceps, *Day*, 147; planifrons, *Day*, 147; platycephalus, *Peters* (fnote), 146; polynema, *Bleek.*, 146; punctillatus, *Rüpp.* (fnote), 147; puntang, *Bleek.*, 146; sadanundio, *Ham. Buch.*, 147; semidoliatus, *Cuv. & Val.*, 147; septemradiatus, *H. B.* (fnote), 148; setosus, *Jerdon* (fnote), 146; sexfasciatus, *Day*, 146; spectabilis, *Günth.* (fnote), 146; stethophthalmus, *Bleek.* (fnote), 146; Stoliczkæ, *Day* (fnote), 147; striatus, *Day*, 147; sublitus, *Cant.* (fnote), 147; tentacularis, *C. & V.*, 147; tredecemradiatus, *H. B.* (fnote), 148; venenatus, *C. & V.* (fnote), 146; ventralis, *C. & V.* (fnote), 147; viridipunctatus, *C. & V.*, 146; zonalternans, *Day*, 147.
- Gogrius, *Day* (fnote), 341; Sykesii, *Day* (fnote), 346.
- Goniocidaris canaliculata, *Agass.*, 65; difference from *Cidaris*, 64.
- Gonothyræa Loveni, attachment of Heliozoa to 303
- Goshawk, new species of 457
- Goura Albertisi, *Salvad.* 318, 503
- Grallæ of New Guinea 320
- Grammatophora barbata 371
- Graucalus angustifrons, *Sharpe*, *n. sp.*, 81, 499; hypoleucus, *Gould*, 81.
- Gromia granulata, *Schulze*, 397; oviformis, *Dujard.*, 397; socialis, *Archer*, 400.
- Günther, Dr. A., notice of two large extinct lizards formerly inhabiting the Mascarene Islands . . . 322
- Gymnetis ramulosa, *Bates* 130
- Gymnocorax senex, *Less.* 501
- Gymnura, incus of 186
- Gyps fulvescens, *Hume*, = *G. fulvus*, 7; fulvus, geog. distrib. of, 4, and map showing range of, Pl. II.; himalayensis, geog. distrib. of, 8, and map showing range of, Pl. III.; hispaniolensis, *Sharpe*, relinquished as a species, = *O. fulvus*, 5; Kolbi, geog. distrib. of, 9; map showing range of, Pl. III.; doubtfully W.-African, 9; indicus, geog. distrib. of, 10; map showing range of, Pl. IV.; occidentalis, remark on, 5; pallescens, *Hume*, = ? *G. indicus* (*Scop.*), 10; Rueppelli, geog. distrib. of, 9; map showing range of, Pl. IV.; rutilans, *Severtzoff*, = *G. fulvus*, 7.
- Habits of ants, bees, and wasps, 217; of hornbills, 156; of the Norwegian lemming, 27, 29.
- Hæckel's opinion of nucleus and cell 262
- Haleyon sanctus, *V. & H.*, 311, 492; Macleayi, *J. & S.*, 492.
- Haliaëtus leucogaster (*Gm.*), 308, 490.
- Haliastur girrenera (*V.*), 308, 490.
- Halichondria panicea, polyzoa on . . 479
- Halicore, ear-bones of 188
- Halitherium, ear-bones of 188
- Halticinae 475
- Hamiltonia, *Swainson* (fnote) . . . 139
- Haplochili 208
- Haplochilus panchax, *Ham. Buch.* 152
- Haploinodotus, *Gill* (fnote) . . . 140
- Hara, *Blyth* (fnote), 341; Buchananani, *Blyth* (fnote), 346; filamentosa, *Blyth* (fnote), 346.
- Harpagones 195
- Hatteria, quadrato-jugal bone of . . 91
- Hedriocystis pellucida 302
- Helicophagus, *Bleek.* (fnote) . . . 342
- Heliphryns variabilis, *Greeff*, 385, 387
- Heliozoa, 263, 287; skeletophora, 292
- Hemiaster, sp., 67, 68, 69, 70; breeding-pouch of, 69, 70.
- Hemibagrus, *Bleek.* (fnote) . . . 341
- Hemisilurus, *Bleek.* (fnote) . . . 342

- | | Page | | Page |
|---|------|---|----------|
| Hesperornis regalis | 92 | Ithyornis dispar, teeth in | 92 |
| Hestia linteata | 196 | Jobi, new species of goshawk from
the island of | 458 |
| Heterophrys myriopoda, <i>Archer</i> ,
392; spinifera, 29; varians,
<i>Schulze</i> , 386. | | Johnius, <i>Bloch</i> (fnote), 140;
anei, <i>Blyth</i> (fnote), 145; cata-
leus, <i>Cuv. & Val.</i> (fnote), 145;
Dussumieri, <i>Cant.</i> (fnote), 145;
Valenciennei, <i>Eyd.</i> (fnote), 145. | |
| Heteropneustes, <i>Müller</i> (fnote) . | 343 | Kangaroos, ear-bones of | 189 |
| Hirundo javanica, <i>Sparrm.</i> | 498 | Kettelia Lowii | 197 |
| Holothuria tremula, direct mode of
development of | 56 | Kryptopterichthys, <i>Bleek.</i> (fnote) | 342 |
| Holothuroidea pedata | 57 | Kryptopterus, <i>Bleek.</i> (fnote) . . . | 342 |
| Homopriion, <i>Hölb.</i> (fnote) | 140 | Labrus tella katchelee, <i>Russ.</i> (fnote.) | 145 |
| Hornbill, description of a new,
155; on the habits of, 156. | | Labyrinthici | 141, 152 |
| Hornera borealis | 484 | —, bony branchial cavity of | 205 |
| Hyalodiscus, 264; rubicundus, 267. | | —, distribution of | 208 |
| Hyalolampe fenestrata | 295 | Labyrinthulea macrocystis, 279;
resting-stage of, 281; vitellina,
279. | |
| Hyalosphenia lata, <i>Schulze</i> | 394 | Laccophilus obscurus, breathing
of | 174 |
| Hydrocissa albirostris, habits in
feeding | 156 | Lacerta viridis | 371 |
| Hydroporus | 162 | Lagenorhynchus, ear-bones of . . . | 187 |
| — elegans, 171; gyllenhalli, 170;
inæqualis, breathing of, 169;
12-pustulatus, 171; pictus, ex-
per. on, 169. | | Lagomys spelæus (fossil lemming?) | 34 |
| Hydrovatus clypealis | 161 | Lagynis baltica, <i>Schultze</i> | 406 |
| — — — experiments as to
breathing | 166 | Lalage humeralis, <i>Gould</i> , 499;
karu (<i>Less. & Garn.</i>), 499. | |
| Hylobates hoolock, partiality for
living birds | 157 | Lampides pseudelpis | 196 |
| Hymenaster nobilis, <i>Wy. Thoms.</i> ,
<i>n. sp.</i> , 74; allied to Pteraster,
76; pellucidus, wide distrib. of,
73. | | Lamprocoecyx lucidus | 310 |
| Hyperia galba, parasitic on Me-
dusæ | 193 | Lasius flavus, 223, 238; niger,
207, 219, 243. | |
| Hyperoodon, ear-bones of | 187 | Laurion corculum | 197 |
| Hyphydrus ovatus, habits of, 162;
experiments on, 167. | | Lecythium | 399 |
| Hypocerinus bethellianus, <i>Wy.</i>
<i>Thoms.</i> , <i>n. sp.</i> , 47, 51, 55; ana-
tomical soft parts of, 54. | | Leiocassis rama, <i>Ham. Buch.</i> | 346 |
| Hypselobagrus, <i>Bleek.</i> (fnote) . . . | 341 | Leiostomus, <i>Cuv. & Val.</i> (fnote) . . | 140 |
| Ianthœnas albigularis, <i>Bp.</i> , 503;
Rawlinsoni, <i>Sharpe</i> , 503. | | Lemmings always go westward,
31; enemies of, 32; living under
the snow, 30; migration and
habits of Norwegian, 27. | |
| Idiocephala nigripennis | 459 | Lemmus norvegicus | 27 |
| Iguana tuberculata, 371; quadrato-
jugal of, 91. | | Lepidoptera coll. by Capt. Pinwill
in Malacca | 115, 196 |
| Ilybius fuliginosus, observ. on the
breathing of | 176 | —, genital organs of | 195 |
| Indian freshwater fishes, geog. dis-
trib. of, 138; families and genera
of, results, 142. | | Lepidosiren, pneumatic duct of,
210; burrowing in mud, 214. | |
| Insectivora, ossicula auditûs of . . | 186 | Leptophrys | 386 |
| Insects, two new Lepidopterous,
from Malacca | 115 | Leucosia affinis, <i>Miers, n. sp.</i> , 107;
brunnea, <i>Miers, n. sp.</i> , 107;
fusco-maculata, <i>Miers, n. sp.</i> ,
107; Perryi, <i>Miers, n. sp.</i> , 107;
pubescens, <i>Miers, n. sp.</i> , 107;
pulcherrima, <i>Miers, n. sp.</i> , 107;
reticulata, <i>Miers, n. sp.</i> , 107;
Whitmeei, <i>Miers, n. sp.</i> , 107. | |
| Irish Gasterostei, F. Day on | 110 | Limopsis, sp. of, obtained by 'Chal-
lenger' exped. | 47 |
| — greyhound pig, note on | 109 | | |
| Isonychus pictus, <i>Sharp, n. sp.</i> . . | 131 | | |

- | Page | Page |
|---|--|
| Lizards, notice of two large extinct, 322; sacral plexus and vertebræ of, 370. | large extinct lizards formerly inhabiting 322 |
| Lobivanellus miles 504 | Mastacembelus, <i>Cuv. & Val.</i> . . . 141 |
| Lobosa, divis. of Thalamophora . 391 | — armatus, <i>Lacép.</i> , 149; Güntheri, <i>Day</i> , 149; malabaricus, <i>Jerd.</i> (ftnote), 149; marmoratus, <i>C. & V.</i> (ftnote), 149; pancalus, <i>H. B.</i> , 149; pentophthalmus, <i>Gronov.</i> (ftnote), 149; ponticerianus, <i>C. & V.</i> (ftnote), 149; punctatus, <i>C. & V.</i> (ftnote), 149; unicolor, <i>C. & V.</i> , 149; zebrinus, <i>Blyth</i> , 149. |
| Locomotion in sarcode organisms . 264 | Mastigamœba aspera 269 |
| Lophogyps occipitalis, geog. distrib. of, 11; map showing range of, Pl. VI. | Matuta granulosa, <i>Miers, sp. n.</i> . . 108 |
| Loriculus vernalis 156 | — lineifera, <i>Miers, sp. n.</i> . . . 108 |
| Lubbock, Sir J., observations on ants, bees, and wasps.—Part IV. 217 | — maculata, <i>Miers, sp. n.</i> . . . 108 |
| Lycomedes Reichei 130 | — obtusifrons, <i>Miers, sp. n.</i> . . . 108 |
| Machæramphus alcinus 308 | — rubro-lineata, <i>Miers, sp. n.</i> . . 108 |
| M'Intosh, Dr. W. C., note on a new example of the Phyllozoicidæ 215 | Matutidæ 108 |
| M'Lachlan, R., on the nymph-stage of the Embidæ, with notes on the habits of the family, &c. . 373 | Medusæ, new species, varieties, and monstrous forms of 190 |
| Macrogathus caudatus, <i>M'Clell.</i> (ftnote), 149; Hamiltonia, <i>M'Clell.</i> (ftnote), 149; undulatus, <i>M'Clell.</i> (ftnote), 149. | Megapodius duperreyi, <i>Less.</i> , 504; tumulus, egg of, 504. |
| Macrones 341 | Megaptera, ear-bones of 187 |
| — armatus, <i>Day</i> , 345; aor, <i>Ham. Buch.</i> , 344; Bleekeri, <i>Day</i> , 345; Blythii, <i>Day</i> , 344; cavasius, <i>H. B.</i> , 345; chryseus, <i>Day</i> , 344; corsula, <i>H. B.</i> , 345; gulis, <i>H. B.</i> , 344; leucophasis, <i>Blyth</i> , 345; malabaricus, <i>Day</i> , 345; microphthalmus, <i>Day</i> , 345; montanus, <i>Jerdon</i> , 345; keletius, <i>C. & V.</i> , 345; oculatus, <i>Jerdon</i> , 345; punctatus, <i>Jerdon</i> , 345; seenghala, <i>Sykes</i> , 344; tengara, <i>H. B.</i> , 345; vittatus, <i>Bloch</i> , 345. | Megathopa candezei, <i>von Harold</i> . 130 |
| Macropterotonus, <i>Lacép.</i> (ftnote). 343 | Melidora collaris, <i>Sharpe, sp. n.</i> . . 313 |
| Macropus, malleus of 189 | Melithreptus albigularis, <i>Gould</i> . 497 |
| Macroscelides, malleus of 186 | Mesoplon, ear-bones of 187 |
| Magosphæra, <i>Hæck.</i> 435 | Meyerina claviformis, species of crustacea living within 506 |
| — how related, 438; planula (ftnote). 436 | Microcometes paludosa 301 |
| Malabar, limits of, for fish fauna (ftnote). 139 | Microcysts of <i>Cienk.</i> 416 |
| Malacca, the butterflies of, 196; new Lepidopterous insects from. 115 | Micreeca flavigaster 498 |
| Malapterus bengalensis, <i>Gray</i> (ftnote) 350 | Microglossum aterritum 491 |
| Malurus alboscapulatus 315, 494 | Microgromia socialis 401 |
| Mammalian ossicula auditûs, morphology of 185 | Micronema, <i>Bleek.</i> (ftnote) 342 |
| Manatus, ear-bones of 188 | Microsicydium, <i>Bleek.</i> (ftnote) . . 140 |
| Manucodia atra (<i>Less.</i>) 317, 500 | Miers, Edward J., notes on the Oxystomatous Crustacea 107 |
| Marsupialia, ossicula auditûs of . 189 | —, on a new genus and species of Crustacea 183 |
| Marsupium of Echinoderms 56 | —, on species of Crustacea living within the Venus's flower-basket (<i>Euplectella</i>) and in Meyerina claviformis 506 |
| Mascarene Islands, notice of two | Migration and habits of Norwegian lemming 27 |

	Page		Page
Monarcha aruensis, 497; carinata, 497; chrysomelas, 497; guttulates, 497; inornata, 497; melanotus, 497; tricolor, 498.		Neophron ginginianus, as to colour of bill of, 16; geog. distrib. of, 15; map showing range of, Pl. VIII.; monachus = Vultur monachus, 2; perenopterus, geog. distrib. of, 13; map showing range of, Pl. VIII.; pileatus, geog. distrib. of, 17; map showing range of, Pl. IX.	
Monas anyli	417	Neptis dindinga, 196; dorelia, 196; gononata, 196; heliodora, 196; leuconota, 196; mamaja, 196; peraka, 196.	
Monera, Haeckel's monog. of	427	Nesocentor chalybeus	81
—, remarks as to position	427	New Guinea, contributions to the ornithology of, Part I., 79; Part II., 305; Part III., 457; Part IV., 486.	
—, tabular systematic arrang. of	435	Nicholson, H. A., and Etheridge, R., on the genus Alveolites	353
Monitor arenaria	371	Norwegian lemming, D. Crotch on 27, 83, 157; R. Collett on, 237.	
Monodon, ear-bones of	187	Noterus sparsus, exper. on	172
Monothalamia	263	Notes on the genus Alveolites and allied forms of Palaeozoic corals	353
— amphistomata	408	Notice of new living Crinoids belonging to the Apiocrinidæ	47
— monostomata	408	Nuclearia simplex, <i>Cienk.</i>	385
Monostomatous thalamophora	391	Nuclei of endosarc in Actinosphaerium, &c.	427
Monotremata, ossicula auditûs of	189	Nucleus, Hertwig and Lesser's account of	264
Morphology of the mammalian ossicula auditûs	185	Numenius cyanopus, 320; uropygialis, <i>Gould</i> , 504.	
Mugil, <i>Artedi</i> , 141; cascasia, <i>H. B.</i> , 150; corsula, <i>Ham. Buch.</i> , 150; dussumieri, <i>Cuv. & Val.</i> (ftnote), 150; hamiltonii, <i>Day</i> , 150; nepalensis, <i>Günth.</i> (ftnote)	150	Nursia abbreviata, 107; plicata, 107; sinuata, <i>Miers, sp. n.</i> , 107.	
Mugilidæ	141, 150	Nyctalemon docile	197
Mus, malleus of	186	Observations on ants, bees, and wasps	217
— islandicus (? sylvaticus)	33	Odontopteryx, ankylosed teeth of, remarks on	92
Myiagra concinna, 498; latirostris, 498.		Œdionyphis, <i>Latr.</i> , 478; Howittii, <i>Baly, n. sp.</i> , 478.	
Myodes lemmus, 27, 327; area of Norwegian habitat, 328; enemies of, 32, 328, 330; epizootic disease of, 330; food of, 328; migrations, directions of, 333; occurrence of great ones, 334; nest of, 328; reproduction, 329; wandering, consequence of strong vitality, 328, 330, 332; schisticolor, <i>Lilljeb.</i> , 329.		Œnops californiana, 24; falklandica, 24; perniger, 23; urubitinga, 24.	
Myogale, malleus of	186	Oligotoma Saundersii, 379; Michaëli (Pl. XXI.), 383.	
Myra carinata, 107; elegans, 107; mamillaris, 107.		Olulaniasis, a parasitic disease of cats	45
Myrmecobius, incus &c. of	189	Olynthia ruficapilla, <i>Burm.</i> , 380; cubana, <i>Hagen</i> , 381.	
Myrmica ruginodis	222	Olyra longicauda, <i>M'Clell.</i> , 347; Burmanica, <i>Day</i> , 348; laticeps, <i>M'Clell.</i> , 348.	
Myxastrum radians	430, 431	Ompok, <i>Lacép.</i> (ftnote), 342; siluroides, <i>Lacép.</i> (ftnote), 348.	
Myxobrachia	434		
Myxodictyum sociale, <i>Haeckel.</i>	432		
Myxomycetæ, as to their position, 417; develop. of, 414; remarks on, 413; sp. ? <i>Cienk.</i> , observ. on, 416.			
Myzomela obscura, <i>Gould</i>	496		
Nandidæ	139, 144		
Nandus marmoratus, <i>Cuv. & Val.</i> (ftnote)	139, 144		
Nangra Buchanani, <i>Day</i> , 351; iridescens, <i>Ham. Buch.</i> , 351; punctata, <i>Day</i> , 351.			
Nematodes of Porpoise	42		
Neobalæna, ear-bones of	187		

Page	Page		
On the geog. distrib. of <i>Accipitres</i>	1	Ornithocheiroidea	96
<i>Onthophagus sharpi</i> , von <i>Harold</i> , 131; <i>tapius</i> , <i>Sharp</i> , <i>n. sp.</i> , 130.		<i>Ornithocheirus</i> , brain-cavity of, 87; comparison, carpals, &c., of, 100; metacarpals of, 101; sections of tooth of, 92.	
<i>Ophichthys boro</i> , exper. with, as to respiration	202	<i>Ornithoptera ruficollis</i>	197
<i>Ophiocephalidæ</i> , 141, 150, 155; accessory gill-cavity of, 205; distribution of, 208; favourite resort of, 209.		<i>Ornithorhynchus</i> , ankylosis of incus and malleus of	189
<i>Ophiocephalus</i> , <i>Block</i> , 141; <i>affinis</i> , <i>Günth.</i> (fnote), 151; <i>amphibius</i> , <i>M'Clell.</i> (fnote), 150; <i>aurantiacus</i> , <i>Ham. Buch.</i> (fnote), 151; <i>aurolineatus</i> , <i>Day</i> (fnote), 150; <i>barca</i> , <i>H. B.</i> , 150; <i>chena</i> , <i>H. B.</i> (fnote), 151; <i>diplogramme</i> , <i>Day</i> (fnote), 151; <i>fuscus</i> , <i>Cuv. & Val.</i> (fnote), 151; <i>gachua</i> , <i>H. B.</i> , 151; <i>gachua</i> , transported in wet handkerchief, 201; <i>grandinosus</i> , <i>Cuv. & Val.</i> (fnote), 150; <i>indicus</i> , <i>M'Clell.</i> (fnote), 151; <i>karrouvei</i> , <i>Lacép.</i> (fnote), 151; <i>Kelaartii</i> , <i>Günth.</i> (fnote), 151; <i>kora molta</i> , <i>Russ.</i> (fnote), 151; <i>latus</i> , <i>H. B.</i> (fnote), 151; <i>leucopunctatus</i> , <i>Sykes</i> , 150; <i>limbatus</i> , <i>Cuv. & Val.</i> (fnote), 151; <i>marginatus</i> , <i>C. & V.</i> (fnote), 151; <i>marilius</i> , <i>H. B.</i> , 150; <i>micropeltes</i> , <i>C. & V.</i> , 151; <i>montanus</i> , <i>M'Clell.</i> (fnote), 151; <i>nigricans</i> , <i>C. & V.</i> (fnote), 150; <i>pseudomarilius</i> , <i>Günth.</i> , 150; <i>punctatus</i> , <i>Block</i> , 151; <i>punctatus</i> from mud, 214; <i>serpentinus</i> , <i>Cuv. & Val.</i> (fnote), 151; <i>stewartii</i> , <i>Playfair</i> , 151; <i>striatus</i> , <i>Block</i> , 151; <i>Theophrasti</i> , <i>Val.</i> (fnote), 150; <i>wrahl</i> , <i>Lacép.</i> (fnote), 151.		<i>Orycteropus</i> , malleus of	188
<i>Ophiocoma didelphys</i> , <i>Wy. Thomson</i> , <i>sp. n.</i>	77	<i>Osphromenus</i> , <i>Lacép.</i> , 141; <i>nobilis</i> , <i>M'Clell.</i> , 153; olfax kept in pots	201
<i>Ophiolepis squamata</i> , devel. of	56	<i>Ossicula auditus</i> , morphology of mammalian	185
<i>Ophiomastix</i> , 335; <i>flaccida</i> , <i>Lyman</i>	337	<i>Otogyps auricularis</i> , geog. distrib. of, 12; map showing range of, Pl. VII.; calvus, geog. distrib. of, 13; map showing range of, Pl. VII.; Dr. Exton's remarks on, 12.	
<i>Ophioscion</i> , <i>Gill</i> (fnote)	140	<i>Otolithus macrophthalmus</i> , <i>Bleek.</i> (fnote)	145
<i>Ophiura texturata</i> , nervous system of	448	<i>Oxystomatous Crustacean</i>	184
<i>Ophiuridæ</i> , description of a new form of, 335; nervous system of, 449.		<i>Pachycephala</i> , <i>sp.</i> , 317; <i>griseiceps</i> , 317.	
<i>Ophiuroidea</i>	76	<i>Pachypora Fischeri</i> , <i>Nich. & Eth.</i> , 362; <i>frondosa</i> , <i>Nich. & Eth.</i> , 362; <i>lamellicornis</i> , <i>Lindström</i> , 362.	
<i>Orca</i> , ear-bones of	187	<i>Pachypterus melanurus</i> , <i>Swains.</i> (fnote), 347; <i>punctatus</i> , <i>Swains.</i> (fnote), 350; <i>trifasciatus</i> , <i>Swains.</i> (fnote), 347.	
<i>Orcella brevirostris</i> , parasites of	36	<i>Pachyrhamphus crassirostris</i>	92
Organisms, recent researches among some of the more simple sarcode, 261; second notice of	385	<i>Palæmon</i> , <i>sp. ?</i>	507
Organization of <i>Ornithosauria</i>	84	<i>Palæosauria</i> of Von Meyer	106
<i>Orgyria turbata</i>	197	<i>Pamphila mæsoïdes</i>	197
<i>Oriolus striatus</i> , <i>Q. & G.</i> 82, 318, 500		<i>Panay</i> , descrip. of new Hornbill from island of	155
		<i>Pandion leucocephalus</i> , <i>Gould</i> , 309, 490	
		<i>Pangasius Buchanani</i> , <i>Cuv. & Val.</i>	347
		<i>Papilio asperi</i>	197
		<i>Paradisea raggiana</i> , <i>Sclater</i> , 318, 487, 500	
		<i>Paralepta</i> , <i>Baly</i> , <i>n. g.</i> , 474; <i>foveicollis</i> , <i>Baly</i> , <i>n. sp.</i> , 475.	
		<i>Parambassis</i> , <i>Bleek.</i> (fnote)	139

- | | Page | | Page |
|---|----------|--|----------|
| Paranandus, <i>Day</i> (fnote) | 139 | Phyllodoce maculata, <i>Johnst.</i> , 215 ; | |
| Parasilurus, <i>Bleek.</i> (fnote) | 342 | <i>citrina</i> , <i>Malmgren</i> , 215. | |
| Parasites of ants, 257 ; of wild animals, difficult to ascertain destructiveness of, 44. | | Phyllodocidæ, on a new example of, 215. | |
| Parasitic crustacean on <i>Aurelia aurita</i> | 193 | Phyllorhina, malleus of | 186 |
| <i>Parra cristata</i> , <i>V.</i> | 505 | <i>Physarium album</i> | 414 |
| <i>Parthenos lilacinus</i> | 196 | Physeter, ear-bones of | 187 |
| Passeres of New Guinea | 315 | Picariæ of N. Guinea | 309 |
| Peach, C. W., observations on British Polyzoa | 479 | Piezorhynchus nitidus, <i>Gould</i> 316, 498 | |
| <i>Pelidnota Belti</i> , <i>Sharp</i> , <i>n. sp.</i> | 132 | <i>Pimelodus anguis</i> , <i>Ham. Buch.</i> (fnote), 347 ; <i>anisurus</i> , <i>M. Clell.</i> (fnote), 345 ; <i>bagarius</i> , <i>H. B.</i> (fnote), 351 ; <i>batasius</i> , <i>H. B.</i> (fnote), 345 ; <i>carcio</i> , <i>H. B.</i> (fnote), 345 ; <i>carnaticus</i> , <i>Jerdon</i> (fnote), 351 ; <i>chandramara</i> , <i>H. B.</i> (fnote), 346 ; <i>djambal</i> , <i>Bleeker</i> (fnote), 347 ; <i>gagata</i> , <i>H. B.</i> (fnote), 350 ; <i>indicus</i> , <i>M. Clell.</i> (fnote), 345 ; <i>menoda</i> , <i>H. B.</i> (fnote), 345 ; <i>nangra</i> , <i>H. B.</i> (fnote), 351 ; <i>pangasius</i> , <i>Cuv. & Val.</i> (fnote), 347 ; <i>platespogon</i> , <i>Val.</i> (fnote), 351 ; <i>rita</i> , <i>H. B.</i> (fnote), 346 ; <i>seengteo</i> , <i>Sykes</i> (fnote), 345 ; <i>silondia</i> , <i>H. B.</i> (fnote), 350 ; <i>urua</i> , <i>H. B.</i> (fnote), 347. | |
| <i>Pelomyxa palustris</i> | 271-273 | <i>Pinaciophora fluviatilis</i> , <i>Greeff.</i> | 297 |
| <i>Pelobius Hermanni</i> , breathing of, 161, 163 ; experiments, summary of, 165. | | <i>Pinacocystis rubicunda</i> | 297 |
| Peneidæ, sp. of | 506 | <i>Pitta novæ guineæ</i> , <i>Müll. & Schleg.</i> | 315, 494 |
| Pentacrinus, undescribed 'Challenger' expedition | 47 | <i>Plagiophrys</i> , <i>Clap. & Lach.</i> | 399 |
| <i>Pentacrinus asteria</i> , 454 ; <i>briareus</i> , 454 ; <i>Mülleri</i> , 455 ; <i>subangularis</i> , 454 ; <i>Wyville-Thomsoni</i> , 455. | | <i>Plagioscion</i> , <i>Gill</i> (fnote) | 140 |
| <i>Perca sari-kullah</i> , <i>Russ.</i> (fnote) | 145 | <i>Plakopus ruber</i> | 270 |
| Percidæ | 142, 139 | <i>Platanista gangetica</i> , parasites of | 35 |
| <i>Periophthalmus</i> , <i>Schn.</i> , 140 ; <i>argenteolineatus</i> , <i>C. & V.</i> (fnote), 148 ; <i>dipus</i> , <i>Bleek.</i> (fnote), 148 ; <i>Freycineti</i> , <i>Cuv. & Val.</i> (fnote), 148 ; <i>fuscatus</i> , <i>Blyth</i> (fnote), 148 ; <i>kalolo</i> , <i>Less.</i> (fnote), 148 ; <i>koelreuteri</i> , <i>Pallas</i> , 148 ; <i>modestus</i> , <i>Cant.</i> (fnote), 148 ; <i>papilio</i> , <i>Bl. Schn.</i> (fnote), 148 ; <i>Schlosseri</i> , <i>Pallas</i> , 148. | | —, malleus of | 188 |
| <i>Petrodromus</i> , malleus of | 186 | <i>Platyarthrus Hoffmannseggii</i> | 257 |
| <i>Phagorus</i> , <i>M. Clell.</i> (fnote) | 343 | <i>Platycephala</i> , <i>Baly</i> , <i>n. g.</i> , 475 ; <i>eximia</i> , <i>Baly</i> , <i>n. sp.</i> , 476. | |
| <i>Phalacrocorax leucogaster</i> , <i>Gould</i> | 505 | <i>Platycrinus</i> , Palæozoic genus allied to living <i>Hyoerinus bethellianus</i> | 51 |
| <i>Phalacronotus</i> , <i>Bleeker</i> (fnote) | 342 | <i>Platyphora</i> , <i>Verral</i> , <i>n. gen.</i> , 259 ; <i>Lubbockii</i> , <i>n. sp.</i> , 260. | |
| — <i>siluroides</i> , <i>Bleeker</i> (fnote) | 348 | <i>Platystacus chaca</i> , <i>H. B.</i> (fnote) | 349 |
| <i>Phalangogonia obesa</i> , <i>Burm.</i> , 129 ; <i>sperata</i> , <i>Sharp</i> , <i>n. sp.</i> , 134 ; <i>stipes</i> , <i>Sharp</i> , <i>n. sp.</i> , 134. | | <i>Plesioneura asmara</i> , 197 ; <i>Pinwillii</i> , 197. | |
| <i>Phanogenia</i> , <i>Lovén</i> | 442 | <i>Pleurophrys sphaerica</i> , <i>Clap. & Lach.</i> , 407 ; <i>amphitrematoides</i> , <i>Arch.</i> , 407. | |
| <i>Phascogale</i> , incus and stapes of | 189 | <i>Plexus</i> , sacral, of Lizards | 370 |
| <i>Phascocarctos</i> , malleus of | 189 | <i>Plotosus arab</i> | 338 |
| <i>Phelsuma seychellense</i> | 325 | <i>Plusiotis aurora</i> , var. | 130 |
| <i>Phileurus nasicornis</i> , <i>Burm.</i> | 130 | <i>Plyctolophus triton</i> (<i>Temm.</i>) | 490 |
| <i>Phlogœnas jobiensis</i> | 310 | <i>Pneumatic duct</i> of fishes | 210 |
| <i>Phocœna</i> , ear-bones of | 187 | <i>Podargus papuensis</i> , <i>Q. & G.</i> | 493 |
| <i>Phonygamia Jamesii</i> , <i>Sharpe</i> , <i>n. sp.</i> , 500 ; <i>keraudreni</i> , 500. | | <i>Podischnus terisander</i> , 130 ; <i>agenor</i> , 130. | |
| <i>Phora formicarum</i> , <i>Verral</i> , <i>n. sp.</i> | 258 | <i>Polyacanthus</i> , <i>Cuv. & Val.</i> , 141 ; | |
| Phoridæ, new genus and species of, 258 ; parasitic on ants, 258. | | | |
| <i>Phractocephalus gogra</i> , <i>Sykes</i> (fnote), 346 ; <i>kuternee</i> , <i>Sykes</i> (fnote), 346. | | | |

- | | Page | | Page |
|---|----------|---|----------|
| cupanus, <i>Cuv. & Val.</i> , 152; signatus, <i>Günth.</i> , 153. | | Pterocryptes, <i>Peters</i> (fnote) . . . | 342 |
| Polyachus marginicollis | 463 | Pterodactylus, bones of skull, comparisons of, 90, 91; cerebral cavity of, 87; hind limbs of, 104; how grouped, 84; Kochii, patagial membranes of, 103; longirostris, brain of, 87; longirostris, comparison of bones of skull of, 90; longirostris, vertebral column of, 94; scolopaceps, skull of, 90. | |
| Polycystina, remarks on | 288 | Pteropus, malleus of low type . . . | 186 |
| Polyergus rufescens (Pl. XVII.), 245; slaves of, 252. | | Ptilonopus aurantiifrons, 503; coronulatus, 320, 503; iozonus, 503. | |
| Polypterus | 209 | Ptilorhis alberti, 501; magnifica, 501. | |
| Polystomella striatopunctata, nucleus in | 411 | Ptilotis analoga, <i>Reichenb.</i> | 497 |
| Polyzoa, observations on British . . . | 479 | Pyxidicula operculata, <i>Ehrenb.</i> . . . | 394 |
| Pompholyxophrys punicea, <i>Archer</i> | 295 | | |
| Porphyrio melanopterus | 320, 505 | Quadrula symmetrica, <i>Schulze</i> . . . | 395 |
| Poteriocrinus | 53 | Quinqueloculina fusca, <i>Brady</i> ; nucleus in | 411 |
| President's Anniversary Address for 1876, 261; for 1877, 385. | | | |
| Priodon, ear-bones of | 288 | Radiolaria, Cienk. on Collosphæra, 426; Haeckel's observ. on structure of, 426; remarks on, 288. | |
| Priopsis argyrozona, <i>Cuv. & Val.</i> (fnote) | 143 | Rallina plumbeiventris, <i>Wall.</i> | 505 |
| Pristolepis, <i>Jerdon</i> , 139, 144; fasciatus, <i>Bleek.</i> , 144; malabaricus, <i>Günth.</i> , 144; marginatus, <i>Jerdon</i> , 144. | | Rama, <i>Bleek.</i> (fnote), 341; Buchanani, <i>Bleek.</i> (fnote), 346. | |
| Prosthecosacter convolutus, 42; inflexus, 42. | | Raphidiophrys elegans, 296; viridis, <i>Archer</i> , 297. | |
| Protamoeba primitiva | 433 | Rectes ferruginea, <i>Bp.</i> | 317, 499 |
| Protista, Haeckel's view of | 261 | Rhamphorhynchus Bucklandi, coracoid of, 98; Gemmingi, comparison of skull of, 90. | |
| Protogenes primordialis | 428 | Rhinogryphus aura, geog. distrib. of, 21; map showing range of, Pl. VI.; californianus, geog. distrib. of, 24; map showing range of, Pl. VII.; falklandicus, geog. distrib. of, 24; map showing range of, Pl. VI.; perniger, geog. distrib. of, 23; urubitinga, geog. distrib. of, 24; map showing range of, Pl. VII. | |
| Protomonas Huxleyi, 418; amyli, 418. | | Rhinoscira, <i>Gill</i> (fnote) | 140 |
| Protomyxa aurantiaca | 428 | Rhipidura gularis, <i>Müll.</i> | 498 |
| Pseudambassis, <i>Bleek.</i> (fnote) | 139 | Rhizoerinus lofotensis, <i>Sars</i> | 54 |
| Pseudecheneis sulcatus, <i>McClell.</i> . . . | 352 | Rhizopoda, Hertwig and Lesser's <i>versus</i> Carpenter's definition of . . . | 390 |
| Pseudembryo of Echinoderms, remark on | 55 | —, Schulze's hypothetical descent of | 412 |
| Pseudetroplus, <i>Bleek.</i> (fnote) | 141 | Rhombosternus antennatus, <i>Baly</i> , <i>sp. n.</i> , 460; gracilicornis, <i>Baly</i> , <i>sp. n.</i> , 461; sulphuripennis, <i>Suffr. MS.</i> , 459. | |
| Pseudeutropius acutirostris, <i>Day</i> , 347; atherinoides, <i>Bloch</i> , 347; garua, <i>Ham. Buch.</i> , 347; goongwaree, <i>Sykes</i> , 347; longimanus, <i>Günth.</i> (fnote), 347; megalops, <i>Günth.</i> (fnote), 347; Mitchellii, <i>Günth.</i> (fnote), 347; murius, <i>H. B.</i> , 347; Sykesii, <i>Jerdon</i> , 347; taakree, <i>Sykes</i> , 347. | | Rhopalocera, on the male genital armature of | 195 |
| Pseudobagrus, <i>Bleek.</i> (fnote) | 341 | Rhynchobdella, <i>Cuv. & Val.</i> | 141 |
| Pseudochlamys patella | 393 | | |
| Pseudogyps africanus, geog. distrib. of, 11; map showing range of, Pl. V.; bengalensis, geog. distrib. of, 11; map showing range of, Pl. V. | | | |
| Pseudorca, ear-bones of | 187 | | |
| Pseudosciaena, <i>Bleek.</i> (fnote) | 140 | | |
| Pseudosilurus, <i>Bleek.</i> (fnote) | 342 | | |
| Pseudospongasius, <i>Bleek.</i> (fnote) . . . | 342 | | |
| Psolus ephippifer, <i>Wy. Thomson</i> , <i>sp. n.</i> , 61; operculatus, 61. | | | |
| Pteraster, develop. of, 56; militaris, 73. | | | |

	Page		Page
Rhynchobdella aculeata, <i>Bloch</i> , 149;		<i>Schn.</i> , 145; maculata, <i>Gray &</i>	
aculeata from mud, 214; aral,		<i>Hardw.</i> (ftnote), 145; miles,	
<i>Bl. Schn.</i> (ftnote), 149; ocellata,		<i>Lacép.</i> , 145; nasus, <i>Steind.</i>	
<i>Cuv. & Val.</i> (ftnote), 149; ori-		(ftnote), 145; osseus, <i>Day</i> , 146;	
entalis, <i>Bl. Schn.</i> (ftnote), 149.		semiluctuosa, <i>C. & V.</i> , 145; sina,	
Rhynchobdellidæ	141	<i>C. & V.</i> , 145; Vogleri, <i>Bleek.</i> , 145.	
Rhynchoeyon, malleus of	186	Sciænidæ	145
Rhyparida, <i>Baly</i>	473	Sciænoides asper, <i>Blyth</i> (ftnote) .	145
— maculicollis, <i>Baly, sp. n.</i>	473	Sciænopis, <i>Gill</i> (ftnote)	140
Rhytidoceros ruficollis	314, 493	Sciurus, malleus of	186
Rhytina, malleus and incus of	188	Scrupocellaria ferox, 480; Macan-	
Rita Buchanani, <i>Bleek.</i> , 346; chry-		drei, 480; scruposa, Pl. XXIII.,	
sea, <i>Day</i> , 346; crucigera, <i>Günth.</i>		479.	
(ftnote), 346; hastata, <i>Val.</i> ,		Scythrops novæ-hollandiæ, <i>Lath.</i> .	492
346; pavimentata, <i>Val.</i> , 346.		Seeley, H. G., on the organization	
Rolleston, George, on the prehisto-		of the Ornithosauria	84
ric British Sus	108	Sharp, D., descrip. of new species	
Romanes, G. J., account of new spe-		of beetles (Scarabæidæ) from	
cies, varieties, and monstrous		Central America	129
forms of Medusæ, II.	190	—, observations on the respira-	
Rotalia, observations on	412	tory action of the carnivorous	
Saccobranchus	205, 209	water-beetles (Dytiscidæ)	161
— a true amphibious fish, 211;		Sharpe, R. Bowdler, contributions	
circulation of, 207; fossilis, air-		to the ornithology of New Gui-	
sacs of, 204, 205; microcephalus,		nea, Part I., 79; Pt. II., 305;	
<i>Günth.</i> (ftnote), 349; mi-		Pt. III., 457; Pt. IV., 486.	
cropis, <i>Günth.</i> , 349.		—, description of a new Horn-	
Saccolobium retusum	374	bill from the Island of Panay .	155
Sacral plexus of lizards	370	—, notes on collection of birds	
Salenia, sp., obtained by 'Challen-		from S.E. New Guinea	79
ger' expedition, 47; varispina,		—, on a new species of Goshawk	
47.		from the island of Jobi	457
Sarcode, Dujardin's definition of .	262	—, on the collection of birds	
Sarcode organisms, recent re-		brought by Mr. Octavius C.	
searches among, 261; second		Stone from S.E. New Guinea .	486
notice of, 385.		—, on ornitholog. coll. formed	
Sarcorhamphus æquatorialis of		by Dr. James in S.E. New	
Sharpe, doubtful, =? <i>S. gryphus</i> ,		Guinea and Yule Island	305
18; <i>gryphus</i> , geog. distrib. of,		Sheat-fishes of India, 338; air-	
18; map showing range of,		vessel subservient to hearing,	
Pl. I.		338; resort to muddy waters .	339
Sarsia, monstrosities of	190	Sicydiops, <i>Bleek.</i> (ftnote)	140
Saurocephalus, comparison of tooth-		Sicydium, <i>C. & V.</i> , 140; fasciatum,	
structure of	93	<i>Day</i> , 148; griseum, <i>Day, n. sp.</i>	
Scarabæidæ, new species of, from		(ftnote), 140, 148.	
Central America	129	Sicyopterus, <i>Gill</i> (ftnote)	140
Schilbeichthys, <i>Bleek.</i> (ftnote) .	342	Silonia diaphina, <i>Swains.</i> (ftnote),	
Schilbe pabo, <i>Sykes</i> (ftnote) . . .	348	346; lurida, <i>Swains.</i> (ftnote),	
Sciæna	140	350.	
— albida, <i>Cuv. & Val.</i> , 145;		Silundia gangetica, <i>Cuv. & Val.</i> ,	
aneus, <i>Bloch</i> , 145; axillaris, <i>C.</i>		350; <i>Sykesii, Day</i> , 349.	
<i>& V.</i> , 145; Belengeri, <i>C. & V.</i> ,		Silurichthys, <i>Bleek.</i> (ftnote) . . .	342
145; Bleekeri, <i>Day</i> , 145; ca-		— Berdmorei, <i>Blyth</i> (ftnote) .	349
runa, <i>C. & V.</i> (ftnote), 145; ca-		Siluridæ, geog. distrib. of	338
rutta, <i>C. & V.</i> , 146; coitor, <i>Ham.</i>		Silurodes, <i>Bleek.</i> (ftnote)	342
<i>Buch.</i> , 145; cuja, <i>H. B.</i> , 145;		— resort of, 209; swim-bladder	
diacanthus, <i>Lacép.</i> , 145; glauc-		of, 210; wounds from spines of,	
cus, <i>Day</i> , 146; maculata, <i>Bl.</i>		340.	
		Silurus anastomus, <i>Cuv. & Val.</i>	

Page	Page
(fnote), 348; anguillaris, <i>Russell</i> (fnote), 349; asotus, <i>C. & V.</i> (fnote), 348; batrachus, <i>Bloch</i> (fnote), 349; biserratus, <i>Swains.</i> (fnote), 349; boalis, <i>Ham. Buch.</i> (fnote), 348; canius, <i>H. B.</i> (fnote), 348; checkra, <i>H. B.</i> (fnote), 348; Cuvieri, <i>Gray</i> (fnote), 350; duda, <i>H. B.</i> (fnote), 348; Dukai, <i>Day</i> (fnote), 348; glanis, 270; indicus, <i>C. & V.</i> (fnote), 348; lamghur, <i>Heckel</i> (fnote), 348; laticeps, <i>Swains.</i> (fnote), 349; microcephalus, <i>C. & V.</i> (fnote), 348; Mülleri, <i>Bleek.</i> (fnote), 348; mysoricus, <i>C. & V.</i> (fnote), 348; singio, <i>H. B.</i> (fnote), 349; wallago, <i>C. & V.</i> (fnote), 348.	109; leucomystax, 108; on the prehistoric British, 108; serofa, var. domesticus, 108; serofa, var. ferus, 108; verrucosus, 108, 109.
Sirenia, ossicula auditus of	492
Sisor rhabdophorus, <i>Ham. Buch.</i>	56
Skull of the prehistoric British Sus, Prof. Rolleston on (abstract)	504
Sloths, ear-bones of	Tadorna radjah, <i>Garn.</i> 321, 505
Smith, Edgar A., descript. of <i>Acantharachna mirabilis</i> , a new form of Ophiuridæ	Tagiades calligana, 197; lavata, 197.
Solanocrinus costatus, 454; Jægeri, 454; scrobiculatus, 454.	Talegallus fuscirostris, <i>Salvad.</i> 504
Sorex, malleus of	Talpa, incus of 186
South-east New Guinea, birds of	Tanysiptera galatea, 312; micro-rhyncha, <i>Sharpe, sp. n.</i> , 311, 493; sylvia, <i>Gould</i> , 493.
Sparus, <i>Russ.</i> (fnote)	Tape-worms of Porpoise 42
Sphærophyma, <i>Baly, n. gen.</i> 478	Tatusia, ear-bones of 188
— Simoni, <i>Baly, n. sp.</i> 479	Tegumen of Lepidoptera 195
Sphecotheres flaviventris, 500; Salvadorii, 500.	Terillus Duboulayi, <i>Baly, n. sp.</i> 471
Sponges, Crustacea living within 506	— foveolatus, <i>Baly, n. sp.</i> 469
Spongicola venusta, <i>De Haan</i> , Pl. XXIV. 506	— perplexus, <i>Baly, n. sp.</i> 470
Squatarola helvetica (<i>L.</i>) 320	— squamosus, <i>Baly, n. sp.</i> 470
Stellifer, <i>Cuv.</i> (fnote) 140	— vittatus, <i>Baly, n. sp.</i> 471
Sterna melanauchen, <i>Temm.</i> 505	Thalamophora 386
Sticklebacks, Irish, Francis Day on, 110; 15-spined, Couch's remarks on develop. of (fnote), 111.	—, how distinguished from Heliozoa, 388; perforata and imperforata, 389.
Stone, Mr. O. C., birds collected by 486	Thaumantias crucifera, <i>Romanes, n. sp.</i> , Pl. XV., 190; helicobostrycha, <i>Romanes, n. sp.</i> , Pl. XV., 190.
Strix delicatulus, 49; flammea, 490.	Thaumantis aliris, <i>West. ♂</i> , = <i>T. pseudaliris</i> , 115; <i>pseudaliris, Butl., n. sp.</i> , 115, 196.
Strongylocentrotus runs far south 79	Thick-walled cysts of Cienk. 416
Strongylognathus testaceus (Pl. XVII.) 258	Thomson, Sir C. Wyr., notice of new living crinoids belonging to the Apicocrinidæ 47
Strongylus commutatus, 45; tetracanthus, 45.	—, notice of some peculiarities in the mode of propagation of certain Echinoderms of the southern sea 55
Sus andamanensis, 108, 109; barbatus, 108, 109; cristatus, 108,	Thyridium, sp., closely resembling <i>Antichira gratiosa</i> 134
	Tiaropsis 190
	— indicans, <i>Romanes, n. sp.</i> (Pl. XV.), 194; oligoplocama, <i>Romanes, n. sp.</i> (Pl. XV.), 194; polydiademata, <i>Romanes, n. sp.</i> (Pl. XV.), 194.
	Tigrisoma heliosylos 321
	Todopsis Bonapartii, <i>Gray</i> , 316, 498; cyanocephala, 498.
	Tolypeutes, ear-bones of 188
	Totanus brevipes, <i>Cuv.</i> 504
	Trematode parasites from the Dolphins of the Ganges 35

	Page		Page
Trichogaster, <i>Bloch</i>	141	Venus's flower-basket, species of	
— chuna, <i>Ham. Buch.</i> , 153; fasciatus, <i>Bl. Schn.</i> , 153; in earthen pots, 201; labiosus, <i>Day</i> , 153; lalius, <i>H. B.</i> , 153.		Crustacea living within	506
Trichoglossus massena, <i>Bp.</i> , 80, 309, 491		Vernal, G. H., descript. of genus of Floridæ parasitic on ants	258
Trichopodus bejeus, <i>H. B.</i> (ftnote), 153; colisa, <i>H. B.</i> (ftnote), 153; cotra, <i>H. B.</i> (ftnote), 153; sota, <i>H. B.</i> (ftnote), 153.		Vertebræ, sacral, of lizards	370
Trichopsis, <i>Kner</i> (ftnote)	141	Vulturifulvus Kolbei, = Gyps Rueppelli, 10; monachus, geog. distrib. of, 2; map showing range of, Pl. I.; monachus mentioned by Fraser, = Neophron monachus (<i>Temm.</i>), 2.	
Trichopus, <i>Lacép.</i> (ftnote)	141	Vulturidæ, geog. distrib. of	1
Trinema acinus, union of	398	Walking fish, experiments with	203
Tropidorhynchus novæ-guineæ, <i>S. Müll.</i>	497	Wallago attu, <i>Bloch</i> , 348; <i>Russellii</i> , <i>Bleek.</i> (ftnote), 348.	
Trout, conveyance to India	199	Water-beetles, on the respiratory action of	161
Tupaia, malleus of	186	White, F. Buchanan, on the male genital armature in European Rhopalocera	195
Umbellularia, sp., obtained by 'Chal-lenger' exped.	47	Wombats, ear-bones of	189
Urapteryx marginipennis	197	Yellow cells of Radiolaria contain starch	427
Vampyrella, <i>Cienk.</i> , 419; gomphonematis, 421; spirogyræ, 419.		Yphthima corticaria	196
Ventral fins, stress laid on presence or absence of	110		

END OF THE THIRTEENTH VOLUME.